

University of Zagreb
Faculty of Science
Department of Biology

Maša Frleta - Valić

Population genetic structure of *Paramuricea clavata* (Risso, 1826) in the eastern Adriatic as revealed by microsatellite markers

Graduation Thesis

Zagreb, 2013

This Thesis,
completed at the Institut de Ciències del Mar, CSIC, Barcelona, Spain
under the supervision of Dr. Jean-Baptiste Ledoux
and co-supervision of Prof. dr. Krunoslav Brčić - Kostić,
was submitted for assessment to the
Department of Biology,
Faculty of Science
at the University of Zagreb
in order to acquire the
Master's degree in Ecology and Nature Protection.

Ovaj rad,
izrađen u Institut de Ciències del Mar, CSIC, Barcelona, Španjolska
pod vodstvom dr. sc. Jean-Baptiste Ledoux
i suvodstvom prof. dr. sc. Krunoslava Brčić - Kostića,
predan je na ocjenu
Biološkom odsjeku
Prirodoslovno-matematičkog fakulteta
Sveučilišta u Zagrebu
radi stjecanja zvanja
Magistre ekologije i zaštite prirode.

Acknowledgements

First of all, my gratitude goes to my supervisor Jean-Baptiste Ledoux for all his help, kindness, guidance and numerous advices through all stages of this research.

I would like to warmly thank Joaquim Garrabou for accepting me as an internship student to Instut de Ciències del Mar in his research group, involving me in his projects and for his constant interest, patience, support and help along the way.

I would also like to thank Jordi, Martina, Clara, Maria, Ariadna, Carlos, Stefano and Edu for beautiful working atmosphere and really nice moments in Barcelona.

It was beautiful experience and pleasure working with all of you.

Zahvale

Hvala mom sumentoru, profesoru Krunoslavu Brčić - Kostiću, što me upoznao s populacijskom genetikom i Silviji Kipson na preporuci za praksu na Institut de Ciències del Mar u Barceloni i uzorcima na kojima je ovaj rad izrađen.

Hvala mami i tati što su pretrpili i isfinancirali sva ronjenja, terene, ljetne škole i ine prohtjeve kroz sve ove godine studiranja.

Hvala BIUS-u što je bio moj drugi dom te svim BIUS-ovcima i BIUS-ovkama što su mi učinili ovo razdoblje studiranja najljepšim ikada.

Hvala bivšim predsjednicima Maji, Dini, Vanji i Vedranu što su vjerovali u mene i naučili me svašta nešto, možda životu najviše.

Hvala najboljoj grupi na svijetu, Grupi za biologiju mora ili Marićki, Čupavom, Zubici, Dariji, Sanji, Vanji, Karli, Teni, Katji, Mirni, Tinče, Petri, Sanji, Borisu, Fatuti, Sonji i svim buddyima koji su mene i koje sam ja učila ronjat i koji su širili sreću na terenima.

Hvala Hrvoju što je uvijek bio poput velikog brata i brinuo se za mene, bilo pod morem, na moru ili uz more, i naučio me puno o ribicama i školjkicama, a sad mi čak da i da vozim Krilavog.

Hvala Marićki i Dariji na svim srijedama, petcima i ostalim danima u tjednu.

Hvala „curama s mora“, Sanji, Tinče, Ariani i Martini na nebrojenim kavama, a i one su bome širile veliku sreću po svuda pa i po Barceloni.

Hvala Vedrani, Anji, Inji, Svenu i drugim „molekularcima“, na svim putovanjima, novim godinama, interdisciplinarnim kavama i budalaštinama.

Hvala Damjanu što je najjači.

University of Zagreb
Faculty of Science
Department of Biology

Graduation Thesis

**Population genetic structure of *Paramuricea clavata* (Risso, 1826)
in the eastern Adriatic as revealed by microsatellite markers**

Maša Frleta - Valić

Rooseveltova trg 6, 10000 Zagreb, Croatia

The red gorgonian *Paramuricea clavata* is a long-lived, slow-growing coral that contributes significantly to the biomass and structural complexity of the Mediterranean hard-bottom communities. In this research, the population genetic structure of *P. clavata* in the eastern Adriatic was assessed for the first time. Thirteen locations in four geographical regions along the Adriatic Sea were genotyped using seven microsatellites. Gene diversity, population structure, dispersal patterns and connectivity among populations were analysed using a hierarchical sampling scheme, from local to regional scale. Results showed strong genetic differentiation which was previously never recorded in the population genetic studies of benthic species in the Adriatic. Discussed results should be valuable for creating or improving management plans for *Paramuricea clavata* and other endangered species with similar life-history traits in the Adriatic Sea.

(42 pages, 8 figures, 5 tables, 98 references, original in: English)

Thesis deposited in the Central Biological Library.

Key words: population genetics, red gorgonian, conservation biology, genetic clustering, isolation by distance

Supervisor: dr. Jean-Baptiste Ledoux
Co-supervisor: prof. dr. Krunoslav Brčić - Kostić
Assisting supervisor: dr. Silvija Kipson

Reviewers: prof. dr. Krunoslav Brčić - Kostić
asst. prof. dr. Petar Kružić
asst. prof. dr. Ana Galov

Thesis accepted: 29 May 2013

Sveučilište u Zagrebu
Prirodoslovno - matematički fakultet
Biološki odsjek

Diplomski rad

**Genetička struktura populacija vrste *Paramuricea clavata* (Risso, 1826)
na istočnom Jadranu određena pomoću mikrosatelita**

Maša Frleta - Valić

Rooseveltova trg 6, 10000 Zagreb

Crvena gorgonija, *Paramuricea clavata* je dugoživući, spororastući sesilni koralj koji čini vrlo bitan dio kompleksnog koraligenskog staništa u cijelom Sredozemnom moru. Ovim istraživanjem, po prvi puta je određena genetska raznolikost ove vrste u Jadranu. Trinaest lokacija iz četiri geografske regije duž istočne jadranske obale je genotipizirano pomoću sedam mikrosatelitskih markera. Genetska raznolikost, struktura populacije, uzorci disperzije te povezanost među populacijama su analizirani na lokalnoj i regionalnoj razini. Također, po prvi puta je pokazana jaka genetska diferencijacija kod bentoskih vrsta u Jadranu. Rezultati ovog istraživanja mogu poslužiti kao vrlo vrijedni podaci za uspostavu i unaprijeđenja menadžment planova za vrstu *Paramuricea clavata* te druge Jadranske ugrožene vrste sa sličnim životnim ciklusom i karakteristikama.

(42 stranice, 8 slika, 5 tablica, 98 literaturnih navoda, jezik izvornika: engleski)

Rad je pohranjen u Središnjoj biološkoj knjižnici.

Ključne riječi: populacijska genetika, crvena gorgonija, konzervacijska biologija Jadransko more, mikrosatelitski markeri

Voditelj: dr. sc. Jean-Baptiste Ledoux
Suvoditelj: prof. dr. sc. Krunoslav Brčić - Kostić
Pomoćni voditelj: dr. sc. Silvija Kipson

Ocjenjitelji: prof. dr. sc. Krunoslav Brčić - Kostić
doc. dr. sc. Petar Kružić
doc. dr. sc. Ana Galov

Rad prihvaćen: 29. svibnja 2013.

Table of contents

| | |
|---|-----------|
| 1 INTRODUCTION | 1 |
| 1.1 Unknown but threatened marine diversity | 1 |
| 1.2 Spatial population genetic structure: studying dispersal in sessile organisms | 3 |
| 1.3 Implications for conservation of marine biodiversity | 5 |
| 1.4 <i>Paramuricea clavata</i> as a studied species | 7 |
| 2 THE AIM OF THE REASERCH | 10 |
| 3 MATERIALS AND METHODS | 11 |
| 3.1 Sampling design | 11 |
| 3.2 Microsatellite analysis | 14 |
| 3.3 Microsatellite characteristics and Hardy - Weinberg equilibrium | 17 |
| 3.4 Pairwise genetic differentiation and Isolation by distance | 18 |
| 3.5 Clustering analyses | 19 |
| 4 RESULTS | 20 |
| 4.1 Microsatellite characteristics and Hardy - Weinberg equilibrium | 20 |
| 4.2 Pairwise genetic differentiation and Isolation by distance | 23 |
| 4.3 Clustering analyses | 25 |
| 5 DISCUSSION | 27 |
| 5.1 Genetic diversity of <i>Paramuricea clavata</i> | 27 |
| 5.2 Population relationships and connectivity | 28 |
| 6 CONCLUSION | 30 |
| 6.1 Implications for the conservation of <i>Paramuricea clavata</i> | 30 |
| 7 REFERENCES | 32 |
| 8 APPENDICES | 42 |

1 INTRODUCTION

1.1 Unknown but threatened marine diversity

There are less than 200.000 catalogued marine species living in the ocean and around 2 million more waiting to be discovered (Mora *et al.* 2011). The biggest knowledge gap is the deep-sea diversity where data is extremely scarce (Webb *et al.* 2010). Unfortunately, with the present rate of discoveries of new species and their extinctions, many of them will disappear before being even discovered and evolutionary relationships among them will not be possible to describe and understand (Hendry *et al.* 2010).

Furthermore, it is also demanding to determine the conservation status of already known species in the ocean. Since little is known of many species' distribution or range, a lot of effort is needed to determine whether populations are stable or changing, and if they are threatened or endangered. A research on conservation priorities of tropical coral reefs revealed that up to 53.6% of some taxons have highly restricted ranges, rendering them vulnerable to extinction (Roberts *et al.* 2002). These data show a lot of reasons for serious concern about world's marine biodiversity.

Mediterranean Sea is one of the world's biodiversity hotspots (Myers *et al.* 2000). With 0.82% of the world's ocean area and 0.3% of its volume, it comprises 4 to 18% of the world's marine species. The main reasons for this high diversity are Mediterranean's tormented geological history which caused numerous environmental changes and led to high speciation due to species' adjustment to cold, temperate and sub-tropical conditions co-occurring in the sea (Bianchi & Morri 2000). Additionally, the rate of endemic species in the Mediterranean is remarkably high, with the average of more than 25% (Lejeusne *et al.* 2009).

Although population genetics is becoming an increasingly frequent topic of research in the world, such studies about the Adriatic species are very scarce. Moreover, they are being conducted only on charismatic species like marine mammals (Gaspari *et al.* 2007) or commercially exploited species like fish (Rossi *et al.* 2006; Tinti *et al.* 2002), cephalopods (Garoia *et al.* 2004) and mussels

(Štambuk *et al.* 2013). Other invertebrates, even endangered ones, are being completely cast aside. Studies based on global scale, even the ones on extremely vagile species (Gaspari *et al.* 2007; Rossi *et al.* 2006) show distinction of Adriatic populations from the NW Mediterranean populations. Research by Garoia *et al.* (2004) and Štambuk *et al.* (2013) thus showed both very low genetic differentiation among populations suggesting the occurrence of single genetically homogeneous populations along the whole Adriatic.

According to these studies, it is of great significance to reveal the spatial genetic structure of the species in natural populations. Defining the spatial distribution of genotypes, it might be possible to understand historical events and selection pressures on investigated species (Vekemans & Hardy 2004).

1.2 Spatial population genetic structure: studying dispersal in sessile organisms

Spatial population genetic studies are based on tracing genetic variation (allele frequencies) between populations along a geographical area. They allow insight into various parameters characterizing the biology of populations such as dispersal (Broquet & Petit 2009). In sessile species, such as marine invertebrates as well as plants, genetic dispersal occurs through propagule. In the marine realm, this propagule (i.e. larvae) is carried away by sea currents and plays an important role in connecting spatially disjunct populations (Gaines *et al.* 2003). Since this environment is a three-dimensional and immense space compared to the size of larvae, revealing their route to settlement can be a challenging task (Hellberg *et al.* 2002) and population genetics approaches are useful tool for overcoming this concern (Féral 2002).

Marine populations are usually much less connected than it would be expected based on dispersal capacity of species' propagules (Marshall *et al.* 2010). Connectivity among populations can be affected by various factors, of which the most obvious ones are physical barriers to larval dispersal (Pineda *et al.* 2007). Place of settlement is determined by characteristics of the oceanographic flow field and two biological variables: time larvae spend in the water column before they become developmentally and physiologically capable of settling (precompetency period), and the time over which larvae settle (competency period) (Jackson & Strathmann 1981). Furthermore, natural selection also seems to play an important role in population connectivity which may be reduced by "phenotype-environment mismatches" i.e. reduction of fitness when an organism that is specialized to one environment finds itself in an alternative environment (DeWitt *et al.* 1998). Consequently, oceanographic barriers and biological characteristics of marine invertebrates can affect gene flows among populations and thereby shape the spatial genetic structure of the species (Baums *et al.* 2006; Ledoux *et al.* 2010).

Until recently, connectivity among marine populations was usually explained by Wright's (1943, 1946) island model, stating all populations are the same size, equally distant one from another, and linked by the equal gene flow, with a

proportion of migrants every generation (Palumbi 2003). Since these conditions are not common in natural environments and there are various other aspects affecting the gene flow, this model is not realistic for populations in nature.

On the other hand, the stepping-stone model assumes that populations that are geographically closer exchange larger amounts of genetic material i.e. genetic differentiation increases with geographical distance. (Kimura & Weiss 1946, Malécot 1950) (Figure 1).




Figure 1 Two common models of population structure: Sewell Wright's island model and Stepping-stone model (Palumbi 2003).

1.3 Implications for conservation of marine biodiversity

Besides species and ecosystems diversity, genetic diversity is the third level of biodiversity that needs particular conservation efforts (McNeely *et al.* 1990) but it is often being neglected. Unfortunately, short-term economic and political interests overcome scientific opinions when creating management and conservation plans (Lande 1988). Additionally, one of the main contributors for huge development in discovering, describing, and organizing biodiversity is the recent improvement of technology and methods. Conservation genetics is an extremely effective tool for identifying the environmental drivers of biodiversity and revealing biodiversity hotspots or areas of endemism. (Hendry *et al.* 2010).

Before the work of Reusch *et al.* (2005) there was no evidence of genetic diversity enhancing ecosystem functionality. Populations of a species are becoming rapidly depleted and consequently the genetic variation is reduced (Frankham 2005a). Depleted ecosystems may not contain the species and genetic diversity necessary to enable them to recover or survive major environmental changes and stresses, such as climate changes, diseases or new predators (Garrabou *et al.* 2009). Furthermore, the decline of one species can lead to the increase or decline of others and thus pose a threat to the stability of the ecosystem. Loss of genetic diversity typically impacts over the long-term, associated with environmental change, while inbreeding depression has an immediate impact on the whole ecosystem (Frankham 2005b).

Since climate changes are not something we can stop, it is necessary to gain insights into species' life history and diversity in order to prevent, or at least minimise, future loss of biodiversity. Additionally, habitat degradation due to human population growth is not helping in species preservation (Lotze *et al.* 2006). A clear understanding of a species' life history and diversity is essential for creating effective and scientifically-based management plans. If both, demography and population genetics are not taken into concern, protecting the habitat may have no effect on ensuring population sustainability (Lande 1988). Habitat restoration and preservation of healthy populations will be possible only by determining the effectiveness of marine protected areas according to dispersal, genetic traits and other species' characteristics (Gaines *et al.* 2003).

Furthermore, population genetic studies can show us how current and past biodiversity patterns have been influenced by past evolutionary processes. Consecutively, these studies can help predict what might happen to biodiversity patterns in the face of future changes in geography and climate (Hendry *et al.* 2010).

An evolutionary perspective is required to evaluate the biological importance of the data. In addition, these data will allow an understanding of past evolutionary events in endangered species, whether they are nonselective, such as bottlenecks or gene flow, or selective, such as detrimental or adaptive mutations (Hedrick 2001).

1.4 *Paramuricea clavata* as a studied species

Paramuricea clavata (Figure 2) is an anthozoan that thrives on steep, vertical or overhanging rocky walls but also on horizontal bottoms if the light level is low enough. It is widespread along the western part of the Mediterranean Sea and in the Adriatic Sea, and sporadically in the Aegean Sea and Atlantic Ocean (Carpine & Grashoff 1975). Populations show patchy distribution and can be found from 5 to 200 meters in depth (Mokhtar-Jamaï *et al.* 2011). As an “ecosystem engineer”, it creates assemblages and it presents a very important part of structurally complex coralligenous habitats (Ballesteros 2006). In addition, it increases the biomass and diversity of the community by providing habitats for other small species.




Figure 2 A colony of *Paramuricea clavata* in the National park Kornati (photo: Hrvoje Čižmek).

Paramuricea clavata is a long-lived species with a low population dynamics. Colony survival rates reach from 63% up to 100% (mean of ~91%) and biggest colonies can be from 60 to 100 years old (Linares *et al.* 2007). Also, its growth rates are very low, from 1.8 to 2.7 cm per year (Coma *et al.* 1998), as well as mean recruitment rates which are around 3% (Linares *et al.* 2007). Compared to other marine invertebrate species, it obtains sexual maturity quite late, between 7 and 13 years of age (Coma *et al.* 1995).

In contrast to tropical gorgonian species where asexual reproduction is common (Bell 1982), *Paramuricea clavata* reproduces only sexually (Coma *et al.* 1995). It is a gonochoric species but there is no sexual dimorphism between male and female colonies. Furthermore, hermaphroditism is very rare. *P. clavata* is iteroparous and reproduces annually. Oogenesis lasts from 13 to 18 months and spawning takes place during June and July (correlated to the sea temperature and the lunar cycle). Since development of new oocytes starts before the ones of the previous year are released, there can be two generations of oocytes found in one polyp. Fertilization occurs internally or on the surface of the mother colony. Following the spawning, first planulas appear after only 48 hours, and may settle down on the substratum in a relatively short time, after spending just a few minutes in the water column (Coma *et al.* 1995).

Health of *Paramuricea clavata* populations is mostly affected by irresponsible SCUBA diving activities, utilization of destructive fishing gear and mass mortality events related to global warming. In areas with a high level of recreational diving, the estimations of the mortality rates of the red gorgonian populations are increasing from 2.7 to 7.4 % per year (Coma *et al.* 2004). During the 2003 heat wave, *P. clavata* was one of the most affected species in the north-western Mediterranean, reaching up to an average of 80% of the colonies impacted (Garrabou *et al.* 2009). Depending on the frequency of extended warm periods or other disturbance events, modelling of near-certain quasi-extinction results in 36 to 84 years (Linares *et al.* 2010). Therefore, it is necessary to gain as much data as possible on the species in order to enhance its conservation. In the eastern Adriatic Sea, the assessment of the *P. clavata* population structure and its conservation status has been initiated recently (Kipson *et al.* 2011).

Previous study on the spatial genetic structure of *Paramuricea clavata* showed strong genetic differentiation at both local and global scale, and a heterozygote deficiency at the sample level. Furthermore, a significant pattern of isolation by distance was observed (Mokhtar-Jamaï *et al.* 2011) at different spatial scale. Likewise, similar studies conducted on other gorgonian species (e.g. *Corallium rubrum*), show strong population genetic structure and limited larval dispersal (Ledoux *et al.* 2010a, Ledoux *et al.* 2010b, Constantini *et al.* 2007). In their study, Mokhtar-Jamaï *et al.* (2011) focused on the north-western Mediterranean populations (Spain, France and west Italy), whereas only one population was sampled at the Croatian coast. Regarding this population, the analysis of STRUCTURE shows that it represents its own cluster with a very low level of admixture with other clusters. This suggests the occurrence of a particular genetic pool in this region. This result combined, with the lack of knowledge regarding the genetic diversity in the Adriatic, call for the development of studies in this particular region of the Mediterranean Basin.

2 THE AIM OF THE RESEARCH

The major goal of this study was to define the spatial distribution of genetic diversity of *Paramuricea clavata* in the eastern part of the Adriatic Sea. To achieve this aim, using a hierarchical sampling scheme (from local to regional scale), two main aspects were analysed:

- i. gene diversity and population structure
- ii. dispersal patterns and connectivity among populations

Obtained results should inform adequate management and conservation of the red gorgonian *Paramuricea clavata* and other endangered species with similar life-history traits in the Adriatic Sea. Data should be taken into consideration for creating scientifically-based No Take Areas and Marine Protected Areas.

3 MATERIALS AND METHODS

3.1 Sampling design

Small tissue fragments of *Paramuricea clavata* colonies were collected at 13 sites (Table 1) following a hierarchical sampling scheme (Figure 4) while SCUBA diving. Locations are distributed in four geographic regions: Senj Archipelago, Kornati Archipelago, Rogoznica Area and Pakleni Islands (Figure 3). Samples were taken at depths from 30 to 50 meters in years 2009 and 2011. At each location, one apical fragment (size: 2 - 3 cm) was taken from around 35 colonies (from 23 to 55) bigger than 20 centimetres in height. Finally, 456 *Paramuricea clavata* tissue fragments were conserved in 95% ethanol and stored at -20°C prior to DNA extraction.




Figure 3 Map of 13 *Paramuricea clavata* sample sites. BAL and BLU are the same location but sampling took place at different depths.

Table 1 Details on sample locations of *Paramuricea clavata*.

| Geographic region | Sample location | Protection status | Population label | Coordinates | | Depth (m) | No. of samples |
|---------------------|-----------------|-----------------------------|------------------|-------------|------------|-----------|----------------|
| | | | | Latitude | Longitude | | |
| Senj Archipelago | Ćutin Mali | unprotected | CUT | 44°43'29"N | 14°29'36"E | 30-40 | 32 |
| Senj Archipelago | Lun | unprotected | LUN | 44°42'15"N | 14°44'02"E | 30-40 | 23 |
| Senj Archipelago | Rt Silo | unprotected | SIL | 44°53'12"N | 14°50'13"E | 30-35 | 30 |
| Senj Archipelago | Vrutak | unprotected | VRU | 44°39'11"N | 14°55'36"E | 38-50 | 48 |
| Rogoznica area | Smokvica | unprotected | SMO | 43°30'38"N | 15°56'32"E | 30-40 | 35 |
| Rogoznica area | Planka | National Ecological Network | PLA | 43°29'36"N | 15°58'09"E | 30-40 | 35 |
| Rogoznica area | Mulo | unprotected | MUL | 43°30'48"N | 15°55'07"E | 30-40 | 35 |
| Kornati Archipelago | Balun | Marine Protected Area | BAL | 43°48'14"N | 15°15'18"E | 30-37 | 31 |
| Kornati Archipelago | Balun | Marine Protected Area | BLU | 43°48'14"N | 15°15'18"E | 35-40 | 39 |
| Kornati Archipelago | Mali Obručan | Marine Protected Area | OBR | 43°50'11"N | 15°13'12"E | 35-40 | 31 |
| Kornati Archipelago | Mana | Marine Protected Area | MAN | 43°48'01"N | 15°15'59"E | 35-40 | 29 |
| Pakleni Islands | Kampanel | Protected landscape | KAM | 43°10'44"N | 16°18'34"E | 30-40 | 55 |
| Pakleni Islands | Lanterna | Protected landscape | LAN | 43°10'05"N | 16°18'38"E | 30-40 | 33 |

Total genomic DNA was extracted from all 456 *P. clavata* individuals using a salting out procedure adjusted from Miller *et al.* (1988) (see Appendix 1 for details).




Figure 4 Hierarchical sampling scheme for one of the four regions sampled across the Eastern Adriatic. Figure modified after Cornell *et al.* (2007).

3.2 Microsatellite analysis

All individuals were genotyped using seven microsatellite loci: Pcla 09, Pcla 10, Pcla 12, Pcla 14, Pcla 17, Pcla 81 (Molecular Ecology Resources Primer Development Consortium *et al.* 2010) and Pcla-a (Agell *et al.* 2009) (Table 2). Markers Pcla 81 and Pcla-a were used for the first time while the rest were already utilized in the study of Mokhtar-Jamaï *et al.* (2011).

All the loci were amplified in 10 µL of PCR solution which contained:

- 20 ng/µL of genomic DNA,
- 1x PCR buffer,
- 2.5 mM MgCl₂,
- 0.5 µM of each primer,
- 0.125 mM dNTPs,
- 0,25 U of GoTaq DNA polymerase (Promega).

The forward primer was labelled with fluorescent dyes (Table 2).

The PCR program was:

- 94°C for 3 min;
- 30 cycles of 94°C for 1 min,
- annealing temperature (60°C for all loci except for Pcla-a 59°C) for 1 min,
- 72°C for 1 min;
- final extension at 72°C for 20 min.

PCR products were analysed on an ABI 3130 Genetic Analyser with GeneScan 600 LIZ internal size standard (Applied Biosystems) at the Plateforme Genome Transcriptome de Bordeaux (France; <http://www4.bordeaux-aquitaine.inra.fr/pgtb>). STRand version 2.2.30 was used for scoring alleles (UC Davis Veterinary Genetics Laboratory) (Figure 5).

Table 2 Primer sequences, PCR conditions and genetic characteristics of seven *Paramuricea clavata* microsatellites.

| Locus name | Primer sequence (5'-3') | Repeat motif | Number of cycles | T _a (°C) | Allele size range (bp) | Genbank accession number |
|------------|--|---------------------|------------------|---------------------|------------------------|--------------------------|
| Pcla 09 | F: 6-FAM_CCTGCTTTTCATGATTTTGC R: CTGTCTAATGTGGCACTGAATG | (CA) ₂₁ | 30 | 60 | 91-126 | GU386261 |
| Pcla 10 | F: 6-FAM_TCGCGTGA CTGCTAACTCTC R: TGTATTGCAGCTGGGATGTG | (GT) ₂₃ | 30 | 60 | 195-237 | GU386262 |
| Pcla 12 | F: VIC_GAAATCGACAATAACACGTTTCG R: TTATCACCTCCGCCAGTTTG | (AC) ₁₅ | 30 | 60 | 100-149 | GU386263 |
| Pcla 14 | F: NED_CTGGGAGTGGGTATCTGAGC R: ACAAGACTTTTCGGAGTGAGG | (CA) ₂₆ | 30 | 60 | 165-215 | GU386264 |
| Pcla17 | F: PET_TCGACGAACTTTGCAGATTG R: CAAATTTCCAAGAAAAGGAAGG | (AC) ₁₄ | 30 | 60 | 170-191 | GU386265 |
| Pcla 81 | F: NED_CCAACAACCGCAAGATAGGT R: TTGCTGCCATATCGTTCATC | (ACA) ₁₀ | 30 | 60 | 175-201 | GU386259 |
| Pcla-a | F: VIC_TGGCAAACATAATTCCACCA R: CAAATGATGCTATTGAAGCCAGT | (GTT) ₉ | 30 | 59 | 158-208 | FJ560968 |

Forward primers are 5' fluorescently labelled with VIC, NED, PET or 6-FAM (Applied Biosystems); T_a annealing temperature; bp = base pair.




Figure 5 STRand version 2.2.30 showing electropherogram of individual PLA16 which is heterozygote for Pcla 10 with alleles 206 and 224 bp peaks. Orange (LIZ) peaks correspond to the size ladder.

3.3 Microsatellite characteristics and Hardy - Weinberg equilibrium

PCR amplification can create DNA fragments that are several repeats shorter than the actual allele. These fragments may be seen as stutter bands (Figure 5) and can be misleading while analysing samples (Ewen *et al.* 2000). Also, as a result of deviations from Hardy-Weinberg proportions, large allele dropouts are possible to occur. Preferential amplification of shorter alleles (Wattier *et al.* 1998) can result in an apparent deficiency in heterozygotes. Consequently, MICRO-CHECKER version 2.2.3 (Van Oosterhout *et al.* 2004) was used to test for scoring errors due to stutters and large allele dropout.

In addition, GIMLET version 1.3.3 was used for identification of identical genotypes that could result from mishandling during sampling processes or PCR preparation (Valière 2002). There is a possibility that same colony might have been sampled twice or that DNA of the same specimen has been replicated twice. Frequencies of null alleles were estimated for each locus and population following the Expectation Maximization (EM) algorithm (Dempster *et al.* 1977) using FreeNA software (Chapuis & Estoup 2007).

Total number of alleles per locus, observed heterozygosity (H_o) and gene diversity i.e. expected heterozygosity (H_e) (Nei 1973) per locus and f estimator of F_{IS} (Weir & Cockerham 1984) were computed with FSTAT version 2.9.3 (Goudet 2001). Observed heterozygosity (H_o) and gene diversity (H_e) (Nei 1973) per population were calculated using GENETIX version 4.05 (Belkhir *et al.* 2004).

Score test for heterozygote deficiency was conducted for each sample with GENEPOP version 4.1.4 (Rousset 2008) in order to test departure from panmixia. Significance of the result was computed by Markov Chain (MC) algorithm (Guo & Thompson 1992; Raymond & Rousset 1995) using default parameters. Null hypothesis of linkage equilibrium was tested for each pair of loci in each sample using 1000 permutations with GENETIX.

3.4 Pairwise genetic differentiation and Isolation by distance

F_{ST} for all populations and all population pairs was computed using Weir & Cockerham's estimator of F_{ST} (θ) in GENEPOP version 4.1.4 (Rousset 2008). Also, pairwise genotyping differentiation between samples was tested with the G test based on default parameters for the Markov chain algorithm (Guo & Thompson 1992).

Isolation by distance (IBD - populations are connected by continuous migration but the rate of gene flow is greatest between neighbouring populations) pattern was analysed through the correlation of genetic and geographic distances between populations (Rousset 1997). Geographic distances in meters were measured among locations using GOOGLE EARTH version 7.0 (<http://earth.google.com>) considering shortest path between each population but avoiding geographical barriers like islands and islets. Also, surface currents were taken into account when determining this route. In GENEPOP, linear regression was made on the genetic distance computed as $F_{ST}/(1-F_{ST})$ over logarithm of geographic distances, as recommended for two-dimensional model (Rousset 1997). The significance of the correlation between two distance matrices was tested by the Mantel test (1967) using 10000 permutations.

AMOVA tests i.e. analyses of molecular variance (Weir & Cockerham 1984, Excoffier *et al.* 1992) were computed using ARLEQUIN version 3.5.1.3 (Excoffier *et al.* 2005) in order to quantify genetic variation among groups (F_{CT}), among populations within groups (F_{SC}) and within populations (F_{ST}). Significance tests were conducted using 1000 permutations. Groups tested with AMOVA were determined by STRUCTURE version 2.2 (Pritchard *et al.* 2000; see below).

3.5 Clustering analyses

The whole dataset was submitted under the Bayesian analysis setup in STRUCTURE version 2.2 (Pritchard *et al.* 2000) in order to evaluate the number of genetic clusters (K) from the individuals' genotypes dataset without any information on their geographical locations or assumptions on any kind of population boundaries (Guillot *et al.* 2005). In computation, it regards admixture, correlated allele frequencies between clusters and uses the recessive allele option to deal with null alleles (Falush *et al.* 2003, 2007). Ten independent runs were performed for each K (varying from 1 to 10) with a burn-in period of 200 000 followed by 500 000 iterations. These values were sufficient to reach stability in parameter estimations. A 'true' K value was then determined based on the logarithm of the likelihood of observing the data - $\ln P(D)$ as a function of genetic cluster - K (Waples & Gaggiotti 2006). The value capturing the major structure in the data is the K value (Pritchard *et al.* 2007).

CLUMPP version 1.1 (Jakobsson & Rosenberg 2007) was used to average the assignment scores over the 10 runs. Lastly, the graphical scheme of results was made using DISTRUCT version 1.1 (Rosenberg 2004).

For multiple tests, significance levels were corrected using a false discovery rate (FDR) correction (Benjamini & Hochberg 1995).

4 RESULTS

4.1 Microsatellite characteristics and Hardy - Weinberg equilibrium

According to MICRO-CHECKER no evidence of scoring errors due to stuttering or large allele dropout was found in the whole data set. In all 13 populations, all loci were polymorphic ($P < 0.01$).

Total number of alleles varied from 5 for Pcla 17 to 29 for Pcla 12, with a mean of 17 alleles per locus. Mean estimated null allele frequencies by locus varied from 0.008 for Pcla 81 to 0.076 for Pcla-a, with a mean value of 0.024 per locus (Appendix 2). No global significant linkage disequilibrium (LD) among loci was observed over all samples (all $P > 0.05$ after FDR correction). Observed heterozygosity ranged between 0.462 for Pcla-a and 0.799 for Pcla 81 (mean of 0.692). Gene diversity varied from 0.484 for Pcla 17 to 0.813 for Pcla 81 (mean of 0.715). Mean H_e observed in MPAs is a bit higher (0.73) than in non protected areas (0.673) but not significantly. Weir & Cockerham's estimator of F_{IS} ranged from -0.107 for Pcla 17 to 0.187 for Pcla-a, with a mean value of 0.035 (Table 3).

Departure from panmixia varied between loci and samples. Heterozygote deficiency was found for two microsatellites, Pcla 10 and Pcla-a and in two populations, LAN and VRU, both with significance of $P < 0.001$ (Appendix 2).

Mean estimated null allele frequencies for populations varied between 0.007 for LUN and 0.057 for VRU with a mean value of 0.024 (Appendix 3). LD among pairs of loci was observed only in LAN population for one pair of loci and in OBR population for two pairs of loci. Observed heterozygosity ranged between 0.616 for VRU and 0.786 for KAM (mean of 0.692). Gene diversity varied from 0.610 for SIL to 0.788 for KAM (mean of 0.704). Multilocus values of Weir & Cockerham's estimator of F_{IS} ranged from -0.042 for SIL to 0.109 for VRU, with a mean value of 0.032 per population. All values were significant at the 0.05 level except VRU population (Table 4).

Table 3 Genetic characteristics of seven *Paramuricea clavata* microsatellites used in this research.

| Locus name | Na | Ho | He | <i>f</i> | r |
|------------|----|-------|-------|---------------|-------|
| Pcla 09 | 20 | 0.796 | 0.785 | -0.015 | 0.006 |
| Pcla 10 | 21 | 0.726 | 0.784 | 0.075 | 0.025 |
| Pcla 12 | 29 | 0.770 | 0.790 | 0.019 | 0.019 |
| Pcla 14 | 17 | 0.753 | 0.790 | 0.049 | 0.024 |
| Pcla17 | 5 | 0.537 | 0.484 | -0.107 | 0.014 |
| Pcla 81 | 19 | 0.799 | 0.813 | 0.025 | 0.008 |
| Pcla-a | 11 | 0.462 | 0.560 | 0.187 | 0.076 |
| mean | 17 | 0.692 | 0.715 | 0.035 | 0.024 |

Na - number of alleles, Ho - observed heterozygosity, He - gene diversity i.e. expected heterozygosity (Nei 1973), *f* - Weir & Cockerham (1984) estimator of F_{IS} , r - null allele frequency. Presented values correspond to the mean value over all samples. Values in bold are significant at the 0.05 level after FDR correction.

Table 4 Measures and estimations of genetic diversity for 13 populations of *Paramuricea clavata* based on seven microsatellite loci.

| Population | Ho | He | f | r |
|------------|---------------|---------------|---------------|-------|
| BAL | 0.678 (0.202) | 0.679 (0.194) | 0.017 | 0.017 |
| BLU | 0.690 (0.140) | 0.724 (0.165) | 0.061 | 0.016 |
| CUT | 0.627 (0.239) | 0.640 (0.190) | 0.035 | 0.021 |
| KAM | 0.786 (0.125) | 0.788 (0.144) | 0.013 | 0.021 |
| LAN | 0.707 (0.194) | 0.771 (0.132) | 0.100 | 0.050 |
| LUN | 0.645 (0.076) | 0.626 (0.126) | -0.008 | 0.007 |
| MAN | 0.732 (0.124) | 0.717 (0.134) | -0.003 | 0.009 |
| MUL | 0.713 (0.189) | 0.748 (0.108) | 0.062 | 0.034 |
| OBR | 0.727 (0.170) | 0.731 (0.165) | 0.023 | 0.014 |
| PLA | 0.694 (0.185) | 0.702 (0.169) | 0.025 | 0.029 |
| SIL | 0.646 (0.266) | 0.610 (0.244) | -0.042 | 0.023 |
| SMO | 0.733 (0.129) | 0.732 (0.115) | 0.014 | 0.020 |
| VRU | 0.616 (0.224) | 0.683 (0.127) | 0.109 | 0.057 |
| mean | 0.692 (0.174) | 0.704 (0.155) | 0.032 | 0.024 |

Ho - observed heterozygosity, He - gene diversity (Nei 1973), f - Weir & Cockerham (1984) estimator of F_{IS} , r - null allele frequency. Presented values correspond to the mean value over all samples. Standard deviations are in brackets. Values in bold are significant at the 0.05 level. Shaded rows are populations in MPAs.

4.2 Pairwise genetic differentiation and Isolation by distance

Pairwise F_{ST} values among all pairs of populations were scattered from 0.0008 between BAL and BLU to 0.178 between BAL and SIL, with a mean value of 0.097 (Appendix 4). After FDR correction, all pairwise comparisons showed highly significant genotypic differentiation observed among all populations.

Presence of the Isolation by distance model of gene flow was confirmed by high significance ($P < 0.001$) of the correlation between $F_{ST}/(1-F_{ST})$ and logarithm of geographic distances (Figure 6).




Figure 6 Linear regression of the genetic distance computed as $F_{ST}/(1-F_{ST})$ over logarithm of geographic distances (m).

The AMOVA showed genetic variation among four clusters of 7.58%, among populations within clusters of 5.39% and within populations 87.03%. Also, high significance of results was observed (Table 5; all $p < 0.001$).

Table 5 AMOVA test - results of the analysis of molecular variance within populations of *Paramuricea clavata*, among its populations within clusters and among clusters. Grouping was based on four clusters defined by STRUCTURE (K=4).

| source of variation | d. f. | variation | P value |
|-----------------------------------|-------|-----------|---------|
| among clusters | 3 | 7.58 % | < 0.001 |
| among populations within clusters | 9 | 5.39 % | < 0.001 |
| within populations | 443 | 87.03 % | < 0.001 |

4.3 Clustering analyses

Using STRUCTURE, analyses made for K spanning from 2 to 4 presented new clusters. For K=2, apparent distinction is shown between Senj archipelago and the rest of the regions. For K=3, Kornati area has separated from Pakleni islands and Rogoznica area with OBR population showing a high level of admixture. For K=4, Pakleni islands and Rogoznica area have clear distinction into two separate clusters (Figure 7, Figure 8).




Figure 7 *Paramuricea clavata* population structure computed by STRUCTURE analysis. Every vertical line represents one colony in the population. Each colour shows individual's ratio in one cluster. Colonies were grouped into two, three and four clusters.




Figure 8 Map of 13 *Paramuricea clavata* sample sites. BAL and BLU are the same location but sampling took place at different depths. Lines divide geographical areas selected for hierarchical sampling scheme. Circles correspond to the level of assignment of each population in one of the four clusters defined by STRUCTURE.

5 DISCUSSION

5.1 Gene diversity and population structure

This study was the first one to characterize the genetic diversity of one engineering cnidarian species in the Adriatic Sea. Microsatellites Pcla 81 and Pcla-a were not used before in the research of *Paramuricea clavata* even if they were already in the gene bank. Therefore, they were efficiently developed at the population research for the first time and now complement previously used ones.

Departure from Hardy-Weinberg equilibrium was shown as well as in other marine sessile animals like sponges (Duran *et al.* 2004), ascidians (Dupont *et al.* 2009; Pérez-Portela & Turon 2008) and corals (Mokhtar-Jamaï *et al.* 2011; Ledoux *et al.* 2010a,b; Polato *et al.* 2010). High genetic diversity was comparable to the West-Mediterranean populations of *Paramuricea clavata* (mean H_e by samples: 0.71 vs. 0.74) while heterozygote deficiency and congruently null allele frequencies were much lower than previously described (Mokhtar-Jamaï *et al.* 2011). Still, it is expected for this species to show heterozygote deficiency at some rate due to its reproductive characteristics. Mating between related individuals is very likely because females brood the eggs on their surface and depend on low sperm dispersal (Coma & Lasker 1997a,b) and also larvae usually settle near the mother colony (Coma *et al.* 1995; Linares *et al.* 2008).

At the maximum of Pleistocenic glaciation, the water level in the Mediterranean Sea was more than 100 meters lower than today. Consequently, only the deepest parts of the Adriatic Sea, the Middle Adriatic Pit and the South Adriatic Pit (Appendix 5a) with surrounding areas remained submerged. A possible explanation for high genetic diversity and lower heterozygote deficiency might be a post-glacial re-colonisation of the Adriatic. Beginning with individuals that remained afterwards isolated in the deepest parts of the Adriatic and gradual colonisation of individuals from deeper Eastern Mediterranean (Ruggero *et al.* 2002). Previous STRUCTURE analysis from the study of Mokhtar-Jamaï *et al.* (2011) showed Adriatic population genetically most similar to the Aegean population which confirms gene flow from the east. A mixture of ancient pre-glaciation genes from the Adriatic and genes from the Eastern Mediterranean create a valuable source of genetic material.

5.2 Dispersal patterns and connectivity among populations

Since the eastern Adriatic coast is well-indented and there is a lack of knowledge on hydrodynamic processes of the sea, it is hard to predict the route of propagules to their settlement. Global surface currents flow from the Strait of Otranto in the south to Gulf of Piran in the north along the eastern coast and back to Otranto along the Italian coast (Appendix 5b, Poulain 2001). Moreover, some studies at local scale showed similarities between surface and deep currents (Vilibić *et al.* 2009) but only a few data exist on this topic in the Adriatic. The actual hydrodynamics occurring in the Adriatic are far more complex than described in the literature to date.

Positive and significant correlation between $F_{ST}/(1-F_{ST})$ and logarithm of geographic distances ($R^2 = 0,3613$; $P < 0,001$) suggests a strong pattern of Isolation by distance but with genetic discontinuities. Previous studies on gorgonian species confirmed the occurrence of IBD model of gene flow at Mediterranean scale, but not always on local scales (Mokhtar-Jamaï *et al.* 2011; Ledoux *et al.* 2010b). The capacity to detect IBD pattern depends on the range of distance values and the variance of estimators that are lower at shorter distances (Rousset 1997).

First round of clustering showed clear distinction of Senj Archipelago from all other populations meaning geographical distance is also an important segment of barriers to gene flow of *Paramuricea clavata*. The reason for this outcome may be a low dispersal of larvae i.e. its short swimming phase (Linares *et al.* 2008) and largest geographical distance from all other populations. AMOVA results confirmed strong cluster genetic differentiation showing higher differentiation among clusters than among populations within clusters (7.58% vs. 5.39%) which have not been recorded yet in octocoralian population genetics studies in the Mediterranean (6.0% vs. 7.2% in Mokhtar-Jamaï *et al.* 2011, 2.7% vs. 6.7% in Ledoux *et al.* 2010b). These high rates of differentiation suggest some heterogeneity between populations within the same group (Planes & Fauvelot 2002). Moreover, this pattern in AMOVA results suggests historical long-term isolation between different regions (Nesbø *et al.* 1998).

Possible explanation to Obručan population (OBR, Kornati Archipelago) belonging to the cluster with populations from the Pakleni Islands might be the strong currents at the southern sides of islands in the past when the contact zone between two clusters was around Kornati Archipelago. Unfortunately, there is no evidence of previous genetic diversity of *Paramuricea clavata* in the Adriatic Sea. Other explanation might be the impact of the sampling scheme. In some cases, different sampling schemes can make STRUCTURE detect different number of clusters (Aurelle & Ledoux 2013).

When it comes to the local scale, populations sampled at the same location but at different depths (30 and 40 meters) - BAL and BLU, did not show any barriers in gene flow as it was the case with some populations in the NW Mediterranean due to thermocline (Mokhtar-Jamaï *et al.* 2011). This outcome was expected since spring-summer thermocline in the middle Adriatic is formed down to a depth of 50 meters (Artegiani *et al.* 1996).

No significant differences in gene diversity were observed between populations inside Marine Protected Areas and the ones living in non-protected areas. Marine protected areas included in this study have been declared in 1963 and 1980, so even considering the low dynamics of *Paramuricea clavata*, some differences in gene diversity were expected in the Kornati National Park and protected landscape Pakleni Islands, in respect to the other non-protected investigated regions (Linares *et al.* 2011). Therefore, this data should be considered as a warning and stimulation to create scientifically-based management plans and a revision of MPA policies for other endangered species and habitats as well.

6 CONCLUSION

6.1 Implications for the conservation of *Paramuricea clavata*

The results of this study provide new information that could improve the management of *Paramuricea clavata* together with similar slow-growing sessile animals and the coralligenous habitat. Because of species' slow dynamics and delicate habitat structure, their recovery from anthropogenic and environmental disturbances is measured in decades (Garrabou *et al.* 2001). Additionally, the presence of strong genetic differentiation suggests these disturbances are usually recovered due to self recruitment, and much more rarely due to the inflow of new genetic material. Therefore, it is necessary to start systematic morphological and genetic monitoring of *P. clavata* and similar vulnerable species (Schwartz *et al.* 2007). Another core measure in conservation of this species is a severe control of diving activities on *P. clavata* sites (Linares *et al.* 2011) since high diving activities seriously damage populations and increase their mortality rates (Coma *et al.* 2004). Regarding extinction predictions due to global warming and other environmental effects (Linares & Doak 2010), it must not be permitted that any other disturbances, especially anthropogenic ones, affect the Adriatic populations.

It is necessary to develop a marine reserve network which would be formed based on the *Paramuricea clavata* dispersal abilities assuring connectivity of populations with a special focus on populations showcasing high genetic diversity (Palumbi 2004). The network of genetic connections among populations is the web that maintains the potential for evolution. Rather than a focus on longisolated populations, efforts should be directed at maintaining networks that capture the adaptive diversity within species (Crandall *et al.* 2000). Moreover, concerning the size of the Adriatic and high genetic differentiation of *Paramuricea clavata* populations, it is an extremely important area for their conservation.

Additionally, the transplantation of colonies in suitable habitats could be a useful solution due to restricted dispersal of larvae (Linares *et al.* 2008). Small populations of endangered species which have lost genetic variation might benefit from the introduction of individuals from related populations or even

subspecies for genetic restoration and recovery to normal levels of genetic variation (Hedrick 2001).

Genetic diversity is essential for further evolution in response to environmental changes and finally, survival of the species (Frankham 2005a). Therefore, genetic diversity definitely is one of the three levels of biodiversity that needs huge attention in conservation (McNeely *et al.* 1990).

7 REFERENCES

- Agell G, Rius M, Pascual M (2009) Isolation and characterization of eight polymorphic microsatellite loci for the Mediterranean gorgonian *Paramuricea clavata*. *Conservation Genetics*, **10**, 2025-2027
- Artegiani A, Bregant D, Paschini E, Pinardi N, Raicich F, Russo A (1996) The Adriatic Sea General Circulation. Part I: Air-Sea Interactions and Water Mass Structure. *Journal of Physical Oceanography*, **27**, 1492-1514
- Ballesteros E (2006) Mediterranean coralligenous assemblages: a synthesis of present knowledge. *Oceanography and Marine Biology: An Annual Review*, **44**, 123-195
- Baums IB, Paris CB, Chérubin LM (2006) A bio-oceanographic filter to larval dispersal in a reef-building coral. *Limnology and Oceanography*, **51 (5)**, 1969-1981
- Belkhir K, Bprsa P, Chikhi L, Raufaste N, Bonhomme F (2004) GENETIX 4.05, logiciel sous Windows TM pour la génétique des populations. Laboratoire Génome, Populations, Interactions, CNRS UMR 5000, Université de Montpellier II, Montpellier, France
- Bell G (1982) The masterpiece of nature: the genetics and evolution of sexuality. University of California Press, Berkeley
- Benjamini Y, Hochberg Y (1995) Controlling the false discovery rate - a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society: Series B*, **57**, 289-300
- Bianchi CN & Morri C (2000) Marine biodiversity of the Mediterranean Sea: situation, problems and prospects for future research. *Marine Pollution Bulletin*, **40**, 367-376
- Broquet T, Petit E (2009) Molecular estimation of dispersal for ecology and population genetics. *Annual Review of Ecology, Evolution, and Systematics*, **40**, 193-216

- Carpine C & Grashoff M (1975) Les Gorgonaires de la Méditerranée. *Bulletin de l'Institut Océanographique de Monaco*, **71 (1430)**, 1-140
- Chapuis MP, Estoup A (2007) Microsatellite null alleles and estimation of population differentiation. *Molecular Biology Evolution*, **24**, 621-631
- Coma R, Lasker HR (1997a) Effects of spatial distribution and reproductive biology on *in situ* fertilization rates of a broadcast-spawning invertebrate. *Biological Bulletin*, **193**, 20-29
- Coma R, Lasker HR (1997b) Small-scale heterogeneity of fertilization success in a broadcast spawning octocoral. *Journal of Experimental Marine Biology and Ecology*, **214**, 107-120
- Coma R, Ribes M, Zabala M, Gili JM (1995) Reproduction and cycle of gonadal development in the Mediterranean gorgonian *Paramuricea clavata*. *Marine Ecology Progress Series*, **117**, 173-183
- Coma R, Ribes M, Gili JM, Zabala M (1998) An energetic approach to the study of life-history traits of two modular colonial benthic invertebrates. *Marine Ecology Progress Series*, **162**, 89-103
- Coma R, Polà E, Ribes M, Zabala M (2004) Long-term assessment of temperate octocoral mortality patterns, protected vs. unprotected areas. *Ecological Applications*, **14**, 1466-1478
- Constantini F, Fauvelot C, Abbiati M (2007) Fine-scale genetic structuring in *Corallium rubrum*: evidence of inbreeding and limited larval dispersal. *Marine Ecology Progress Series*, **340**, 109-119
- Cornell HV, Karlson RH, Hughes TP (2007) Scale dependent variation in coral community similarity across sites, islands, and island groups. *Ecology*, **88**, 1707-1715
- Dempster A, Laird N, Rubin D (1977) Maximum likelihood from incomplete data via the EM algorithm. *Journal of the Royal Statistical Society: Series B*, **39**, 1-38
- DeWitt TJ, Sih A, Wilson DS (1998) Costs and limits of phenotypic plasticity. *Trends in Ecology and Evolution*, **13**, 77-81

- Dupont L, Viard F, Dowell MJ, Wood C, Bishop JDD (2009) Fine- and regional-scale genetic structure of the exotic ascidian *Styela clava* (Tunicata) in southwest England, 50 years after its introduction. *Molecular Ecology*, **18**, 442-453
- Duran S, Pascual M, Estoup A, Turon X (2004) Strong population structure in the marine sponge *Crambe crambe* (Poecilosclerida) as revealed by microsatellite markers. *Molecular Ecology*, **13**, 511-522
- Ewen KR, Bahlo M, Treloar SA, Levinson DF, Mowry B, Barlow JW, Foote SJ (2000) Identification and analysis of error types in high-throughput genotyping. *American Journal of Human Genetics*, **67**, 727-736
- Excoffier L, Laval G, Schneider S (2005) Arlequin ver. 3.0: An integrated software package for population genetics data analysis. *Evolutionary Bioinformatics Online*, **1**, 47-50
- Excoffier L, Smouse PE, Quattro JM (1992) Analysis of Molecular Variance Inferred From Metric Distances Among DNA Haplotypes: Application to Human Mitochondrial DNA Restriction Data. *Genetics*, **131**, 479-491
- Falusch D, Stephens M, Pritchard JK (2003) Inference of population structure using multilocus genotype data: linked loci and correlated allele frequencies. *Genetics*, **164**, 1567-1587
- Falusch D, Stephens M, Pritchard JK (2007) Inference of population structure using multilocus genotype data: dominant markers and null alleles. *Molecular Ecology Notes*, **7**, 574-578
- Féral JP (2002) How useful are the genetic markers in attempts to understand and manage marine biodiversity? *Journal of Experimental Marine Biology and Ecology*, **268**, 121-145
- Frankham R (2005a) Ecosystem recovery enhanced by genotypic diversity. *Heredity*, **95**, 183
- Frankham R (2005b) Genetics and extinction. *Biological conservation*, **126**, 131-140

- Gaines SD, Gaylord B, Largier JL (2003) Avoiding current oversights in marine reserve design. *Ecological Applications*, **13**, 32–46
- Garrabou J, Perez T, Sartoretto S, Harmelin JG (2001) Mass mortality event in red coral *Corallium rubrum* populations in the Provence region (France, NW Mediterranean). *Marine Ecological Progress Series*, **217**, 263-272
- Garrabou J, Coma R, Bensoussan N, Bally M, Chevaldonné P, Cigliano M, Diaz D, Harmelin JG, Gambi MC, Kersting DK, Ledoux JB, Lejeusne C, Linares C, Marschal C, Pérez T, Ribes M, Romano JC, Serrano E, Teixido N, Torrents O, Zabala M, Zuberer F, Cerrano C (2009) Mass mortality in Northwestern Mediterranean rocky benthic communities: effects of the 2003 heat wave. *Global Change Biology*, **15**, 1090-1103
- Garoia F, Guarniero I, Ramšak A, Ungaro N, Landi M, Piccinetti C, Mannini P, Tinti F (2004) Microsatellite DNA variation reveals high gene flow and panmictic populations in the Adriatic shared stocks of the European squid and cuttlefish (Cephalopoda). *Heredity*, **93**, 166–174
- Gaspari S, Azzellino A, Airoidi S, Hoelzel AR (2007) Social kin associations and genetic structuring of striped dolphin populations (*Stenella coeruleoalba*) in the Mediterranean Sea. *Molecular Ecology*, **16**, 2922–2933
- Goudet J (2001) FSTAT, a program to estimate and test gene diversities and fixation indices (version 2.9.3). Available from <http://www.unil.ch/izea/software/fstat.html>. Updated from Goudet (1995)
- Guillot G, Estoup A, Mortier F, Cosson JF (2005) A spatial statistical model for landscape genetics. *Genetics*, **170 (3)**, 1261 - 1280
- Guillot G, Leblois R, Coulon A, Frantz AC (2009) Statistical methods in spatial genetics. *Molecular Ecology*, **18**, 4734-4756
- Guo S, Thompson E (1992) Performing the exact test of Hardy-Weinberg proportion for multiple alleles. *Biometrics*, **48**, 361-372
- Hedrick PW (2001) Conservation genetics: where are we now? *Trends in Ecology & Evolution*, **16**, 629-636

- Hellberg ME, Burton RS, Neigel JE, Palumbi SR (2002) Genetic assessment of connectivity among marine populations. *Bulletin of Marine Science*, **70 (1)**, 273-290
- Hendry AP, Lohmann LG, Conti E, Cracraft J, Crandall KA, Faith DP, Häuser C, Joly CA, Kogure K, Larigauderie A, Magallón, Moritz C, Tillier S, Zardoya R, Prieur-Richard AH, Walther BA, Yahara T, Donoghue MJ (2010) Evolutionary biology in biodiversity science, conservation and policy: a call to action. *Evolution*, **64**, 1517-1528
- Jackson GA, Strathmann RR (1981) Larval mortality from offshore mixing as a link between precompetent and competent periods of development. *The American Naturalist*, **118**, 16-26
- Jakobsson M, Rosenberg NA (2007) CLUMPP: a cluster matching and permutation program for dealing with label switching and multimodality in analysis of population structure. *Bioinformatics*, **23**, 1801-1806
- Kimura M, Weiss GH (1964) The stepping stone model of population structure and the decrease of genetic correlation with distance. *Genetics*, **49(4)**, 561
- Kipson S, Garrabou J, Linares C, Cebrian E, Ballesteros E, Čížmek H, Bakran-Petricioli T, Kružić P (2011) Conservation status of the red gorgonian *Paramuricea clavata* (Risso, 1826) populations from the Eastern Adriatic Sea. *Book of Abstracts of the 46th European Marine Biology Symposium (46th EMBS, Rovinj, Hrvatska, 12 - 16 September 2011)*, Ruđer Bošković Institute, Centar za istraživanje mora Rovinj, Rovinj, Croatia, p. 84
- Koehn RK, Newell RIE, Immerman F (1980) Maintenance of an aminopeptidase allele frequency cline by natural selection. *Proceedings of Natural Academy of Sciences (USA)*, **77**, 5385-5389
- Lande R (1988) Genetics and demography in biological conservation. *Science*, **241**, 1455-1460
- Ledoux JB, Garrabou J, Bianchimani O, Drap P, Féral JP, Aurelle D (2010a) Fine-scale genetic structure and inferences on population biology in the threatened Mediterranean red coral, *Corallium rubrum*. *Molecular Ecology*, **19**, 4204-4216

- Ledoux JB, Mokhtar-Jamaï K, Roby C, Féral, Garrabou J, Aurelle D (2010b) Genetic survey of shallow populations of the Mediterranean red coral [(*Corallium rubrum*, Linnaeus, 1758)]: new insights into evolutionary processes shaping nuclear diversity and implications for conservation. *Molecular Ecology*, **19**, 675-690
- Lejeusne C, Chevalodonné P, Pergent-Martini C, Boudouresque CF, Pérez T (2009) Climate change effects on a miniature ocean: the highly diverse, highly impacted Mediterranean Sea. *Trends in ecology and Evolution*, **25 (4)**, 250-260
- Linares C, Doak DF, Coma R, Díaz D, Zabala M (2007) Life history and viability of a long-lived marine invertebrate: the octocoral *Paramuricea clavata*. *Ecology*, **88 (4)**, 918-928
- Linares C, Coma R, Mariani S, Díaz D, Hereu B, Zabala M (2008) Early life history of the Mediterranean gorgonian *Paramuricea clavata*: implications for population dynamics. *Invertebrate Biology*, **127**, 1-11
- Linares C, Doak DF (2010) Forecasting the combined effects of disparate disturbances on the persistence of long-lived gorgonians: a case study of *Paramuricea clavata*. *Marine Ecology Progress Series*, **402**, 59-68
- Linares C, Garrabou J, Hereu B, Díaz D, Marschal C, Sala E, Zabala M (2011) Assessing the effectiveness of marine reserves on unsustainably harvested long-lived sessile invertebrates. *Conservation Biology*, **26**, 88-96
- Lotze HK, Lenihan HS, Bourque BJ, Bradbury RH, Cooke RG, Kay MC, Kidwell SM, Kirby MX, Peterson CH, Jackson JBC (2006) Depletion, degradation, and recovery potential of estuaries and coastal seas. *Science*, **312 (5781)**, 1806-1809
- Malécot G (1950) Quelques schémas probabilistes sur la variabilité des populations naturelles. *Annales de l'Université de Lyon A*, **13**, 37-60.
- Mantel N (1967) The detection of disease clustering and a generalized regression approach. *Cancer Research*, **27**, 209-220

- Marshall DJ, Monro K, Bode M, Keough MJ, Swearer S (2010) Phenotype-environment mismatches reduce connectivity in the sea. *Ecology Letters*, **13**, 128-140
- McNeely J, Miller K, Reid W, Mittermeier R, Werner T (1990) Strategies for conserving biodiversity. *Environment*, **32**, 16-20
- Miller SA, Dykes DD, Polesky HF (1988) A simple salting out procedure for extracting DNA from human nucleated cells. *Nucleic Acids Research*, **16**, 1215
- Mokhtar-Jamaï K, Pascual M, Ledoux JB, Coma R, Féral JP, Garrabou J, Aurelle D (2011) From global to local genetic structuring in the red gorgonian *Paramuricea clavata*: the interplay between oceanographic conditions and limited larval dispersal. *Molecular Ecology*, **20 (16)**, 3291–3305
- Molecular Ecology Resources Primer Development Consortium, Aurelle D, Baker AJ *et al.* (2010) Permanent genetic resources added to the molecular data base 1 February 2010 - 31 March 2010. *Molecular Ecology Resources*, **10**, 751-753. <http://tomato.bio.trinity.edu/manuscripts/10-4/mer-10-0076.pdf>
- Mora C, Tittensor DP, Adl S, Simpson AGB, Worm B (2011) How Many Species Are There on earth and in the Ocean? *PLOS Biology*, **9 (8)**, e1001127
- Myers N, Mittermeier RA, Mittermeier CG, da Fonesca GAB, Kent J (2000) Biodiversity hotspots for conservation priorities. *Nature*, **403**, 853-858
- Nei M (1973) Analysis of gene diversity in subdivided populations. *Proceedings of the National Academy of Sciences the United States of America*, **70**, 3321-3323.
- Nesbø CL, Magnhagen C, Jakobsen KS (1998) Genetic differentiation among stationary and anadromous perch (*Perca fluviatilis*) in the Baltic Sea. *Hereditas*, **129**, 241-249
- Palumbi SR (2003) Population genetics, demographic connectivity, and the design of marine reserves. *Ecological Implications*, **13 (1)**, S146-S158
- Palumbi SR (2004) Marine reserves and ocean neighbourhoods: the spatial scale of marine populations and their management. *Annual Review of Environmental Resources*, **29**, 31-68

- Pérez-Portela R, Turon X (2008) Cryptic divergence and strong population structure in the colonial invertebrate *Pycnoclavella communis* (Ascidacea) inferred from molecular data. *Zoology*, **111**, 163-178
- Planes S, Fauvelot C (2002) Isolation by distance and vicariance drive genetic structure of a coral reef fish in the Pacific ocean. *Evolution*, **56 (2)**, 378-399
- Pineda J, Hare JA, Sponaugle S (2007) Larval transport and dispersal in the coastal ocean and consequences for population connectivity. *Oceanography*, **20**, 22-39
- Polato NR, Conception GT, Toonen RJ, Baums IB (2010) Isolation by distance across the Hawaiian Archipelago in the reef-building coral *Porites lobata*. *Molecular Ecology*, **19**, 4661-4677
- Poulain PM (2001) Adriatic Sea surface circulation as derived from drifter data between 1990 and 1999. *Journal of Marine Systems*, **29**, 3-32
- Pritchard JK, Stephens M, Donnelly P (2000) Inference of population structure using multilocus genotype data. *Genetics*, **155**, 945-959
- Pritchard JK, Wen X, Falush D (2007) Documentation for the STRUCTURE software, Version 2. Chicago <http://pritch.bds.uchicago.edu>
- Raymond M, Rousset F (1995) GENEPOP (ver. 1.2): a population genetics software for exact test and ecumenicism. *Journal of Heredity*, **86**, 248-249
- Reusch TBH, Ehlers A, Hämmerli A, Worm B (2005) Ecosystem recovery after climatic extremes enhanced by genotypic diversity. *Proceedings of the National Academy of Sciences of the United States of America*, **102 (8)**, 2826-2831
- Roberts CM, McClean CJ, Veron JEN, Hawkins JP, Allen GR, McAllister DE, Mittermeier CG, Schueler FW, Spalding M, Wells F, Vynne C, Werner TB (2002) Marine Biodiversity Hotspots and Conservation Priorities for Tropical Reefs. *Science*, **295**, 1280-1284
- Rosenberg NA (2004) DISTRUCT: a program for graphical display of population structure. *Molecular Ecology Notes*, **4**, 137-138

- Rossi AR, Perrone E, Sola L (2006) Genetic structure of gilthead seabream, *Sparus aurata*, in the Central Mediterranean Sea. *Central European Journal of Biology*, **1(4)**, 636–647
- Rousset F (1997) Genetic differentiation and estimation of gene flow from FStatistics under isolation by distance. *Genetics*, **145**, 1219-1228
- Rousset F (2008) Genepop'007: a complete reimplementation of the Genepop software for Windows and Linux. *Molecular Ecology Resources*, **8**, 103-106
- Rugiero MV, Turk R, Procaccini G (2002) Genetic identity and homozygosity in North-Adriatic populations of *Posidonia oceanica*: An ancient, post-glacial clone? *Conservation Genetics*, **3**, 71-74
- Schwartz MK, Luikart G, Waples RS (2007) Genetic monitoring as a promising tool for conservation and management. *Trends in Ecology & Evolution*, **22**, 25-33
- Štambuk A, Šrut M, Šatović Z, Tkalec M, Klobučar GIV (2013) Gene flow vs. pollution pressure: Genetic diversity of *Mytilus galloprovincialis* in eastern Adriatic. *Aquatic toxicology*, **136-137**, 22-31
- Tinti F, Di Nunno C, Guarniero I, Talenti M, Tommasini S, Fabbri E, Piccinetti C (2002) Mitochondrial DNA Sequence Variation Suggests the Lack of Genetic Heterogeneity in the Adriatic and Ionian Stocks of *Sardina pilchardus*. *Marine Biotechnology*, **4**, 163–172
- Valière N (2002) GIMLET: a computer program for analysing genetic individual identification data. *Molecular Ecology Notes*, **2**, 377-379
- Van Oosterhout C, Hutchinson WF, Wills DPM, Shipley P (2004) MICRO-CHECKER: software for identifying and correcting genotyping errors in microsatellite data. *Molecular Ecology Notes*, **4**, 535-538
- Vekemans X & Hardy OJ (2004) New insights from fine-scale spatial genetic structure analyses in plant populations. *Molecular Ecology*, **13**, 921-935
- Vilibić I, Book JW, Beg Paklar B, Orlić M, Dadić V, Tudor M, Martin PJ, Pasarić M, Grbec B, Matic F, Mihanović H, Morović M (2009) West Adriatic coastal water excursions into the East Adriatic. *Journal of Marine Systems*, **78**, S132-S156

- Waples RS, Gaggiotti O (2006) What is a population? An empirical evaluation of some genetic methods for identifying the number of gene pools and their degree of connectivity. *Molecular Ecology*, **15**, 1419-1439
- Wattier R, Engel CR, Saumitou-Laprade P, Valero M (1998) Short allele dominance as a source of heterozygote deficiency at microsatellite loci: experimental evidence at the dinucleotide locus Gv1CT in *Gracilaria gracilis* (Rhodophyta). *Molecular Ecology*, **7**, 1569–1573
- Webb TJ, Vanden Berghe E, O’Dor RK (2010) Biodiversity’s big wet secret: The global distribution of marine biological records reveals chronic under-exploration of the deep pelagic ocean. *PLoS One*, **5 (8)**, e10223
- Weir BS, Cockerham CC (1984) Estimating F-statistics for the analysis of population structure. *Evolution*, **38**, 1358-1370
- Wright S (1943) Isolation by distance. *Genetics*, **28**, 114–138
- Wright S (1946) Isolation by distance under diverse systems of mating. *Genetics*, **31**, 39–59

8 APPENDICES

Appendix 1

DNA extraction protocol: salting out procedure, adjusted from Miller *et al.* (1988)

Appendix 2

Hardy Weinberg test when $H1$ = heterozygote deficit [Score (U) test], computed in GENEPOP v.4.1.4 (Rousset 2008)

Appendix 3

Table of estimated null allele frequencies, computed with FreeNA software using the EM algorithm (Dempster *et al.* 1977)

Appendix 4

Genetic and geographic distance matrices used for IBD analyses, computed in GENEPOP version 4.1.4 (Rousset 2008)

Appendix 5a

Adriatic Sea coastline and topography, included from Artegiani *et al.* (1996)

Appendix 5b

Map of surface mean flow in the Adriatic basin during summer, included from Poulain (2001)

Appendix 1

DNA extraction protocol: salting out procedure, adjusted from Miller *et al.* (1988)

1. For each colony/individual dissect around 10 polyps in a Petri dish with a clean scalpel
2. Incubation few minutes at 55°C to dry the remaining alcohol
3. Add 600 µL of TNES buffer
4. Add 20 µL of proteinase K (20mg/mL) + vortex + centrifuge
5. Incubation at 55°C for 4 to 5 hours
6. Put the digestion product in a new tube (without any remaining tissue)
7. Add 166,7 µL of NaCl (6M) (during this step the proteins will be precipitated)
8. Centrifugation for 15 min at 10000 g or 10,0 rcf or 12,2 rpm
9. Put the supernatant in a new tube
10. Add 780 µL of cold (-20°C) absolute alcohol (ethanol 100%)
11. Add 78 µL of Sodium Acetate (3M, pH=7)
12. Centrifugation for 15 min at 13000 g or 13,0 rcf or 13,9 rpm (DNA precipitation)
13. Remove the alcohol by pipetting
14. Add 780 µL of cold 70% ethanol
15. Centrifugation for 15 min at 13000 g
16. Remove the alcohol
17. Let the tubes dry over night
18. If needed, incubation for few minutes at 55°C to remove the remaining alcohol
19. Add 100 µL of milliQ sterilized H₂O

Appendix 2

Hardy Weinberg test when H1= heterozygote deficit [Score (U) test], computed in GENEPOP v.4.1.4 (Rousset 2008)

Markov chain parameters for all tests:

Dememorization: 10000
 Batches: 20
 Iterations per batch: 5000

Results by population (test multi-locus):

| Population | P-val | S.E. | switches (ave.) |
|--------------|---------------|---------------|-----------------|
| BAL31 | 0.1114 | 0.0145 | 13415.57 |
| BLU40 | 0.0723 | 0.0120 | 16078.57 |
| CUT33 | 0.1362 | 0.0145 | 19176.43 |
| KAM55 | 0.0241 | 0.0092 | 19562.29 |
| LAN35 | 0.0006 | 0.0006 | 17665.43 |
| LUN24 | 0.5061 | 0.0120 | 20515.71 |
| MAN30 | 0.3389 | 0.0306 | 15281.00 |
| MUL35 | 0.0081 | 0.0020 | 15076.14 |
| OBR31 | 0.1339 | 0.0143 | 18384.57 |
| PLA35 | 0.0291 | 0.0055 | 19177.43 |
| SIL30 | 0.2406 | 0.0180 | 17696.43 |
| SMO36 | 0.1012 | 0.0146 | 14405.29 |
| VRU48 | 0.0000 | 0.0000 | 12861.57 |

Results by locus (test multi-population):

| Locus | P-val | S.E. | switches (ave.) |
|---------------|---------------|---------------|-----------------|
| Pcla09 | 0.7398 | 0.0237 | 7438.69 |
| Pcla10 | 0.0002 | 0.0002 | 9332.46 |
| Pcla12 | 0.0109 | 0.0056 | 3614.69 |
| Pcla14 | 0.0078 | 0.0029 | 9688.23 |
| Pcla17 | 0.8911 | 0.0036 | 59898.23 |
| Pcla81 | 0.2332 | 0.0159 | 9234.15 |
| Pclaa | 0.0000 | 0.0000 | 18876.23 |

Appendix 3

Table of estimated null allele frequencies

FreeNA: Estimating null allele frequency using the EM algorithm (Dempster *et al.* 1977)

| Population: | BAL31 | BLU40 | CUT33 | KAM55 | LAN35 | LUN24 | MAN30 | MUL35 | OBR31 | PLA35 | SIL30 | SMO36 | VRU48 |
|-------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Locus: | | | | | | | | | | | | | |
| Pcla09 | 0,0001 | 0,0000 | 0,0000 | 0,0199 | 0,0503 | 0,0000 | 0,0028 | 0,0000 | 0,0000 | 0,0000 | 0,0000 | 0,0000 | 0,0000 |
| Pcla10 | 0,0695 | 0,0216 | 0,0014 | 0,0162 | 0,0000 | 0,0005 | 0,0279 | 0,0000 | 0,0090 | 0,0252 | 0,0000 | 0,0852 | 0,0654 |
| Pcla12 | 0,0000 | 0,0000 | 0,0000 | 0,0000 | 0,0572 | 0,0000 | 0,0075 | 0,0284 | 0,0000 | 0,0587 | 0,0581 | 0,0073 | 0,0286 |
| Pcla14 | 0,0130 | 0,0544 | 0,0000 | 0,0274 | 0,0000 | 0,0459 | 0,0000 | 0,0713 | 0,0679 | 0,0326 | 0,0000 | 0,0000 | 0,0000 |
| Pcla17 | 0,0376 | 0,0005 | 0,0001 | 0,0000 | 0,0000 | 0,0000 | 0,0000 | 0,1398 | 0,0046 | 0,0000 | 0,0000 | 0,0000 | 0,0000 |
| Pcla81 | 0,0000 | 0,0352 | 0,0000 | 0,0000 | 0,0084 | 0,0000 | 0,0000 | 0,0000 | 0,0148 | 0,0000 | 0,0000 | 0,0000 | 0,0474 |
| Pclaa | 0,0000 | 0,0000 | 0,1456 | 0,0827 | 0,2357 | 0,0000 | 0,0280 | 0,0000 | 0,0000 | 0,0870 | 0,1014 | 0,0491 | 0,2563 |

Appendix 4

Genetic and geographic distance matrices used for IBD analyses computed in GENEPOP v.4.1.4 (Rousset 2008)

Fst/(1-Fst) estimates:

| | BAL | BLU | CUT | KAM | LAN | LUN | MAN | MUL | OBR | PLA | SIL | SMO | VRU |
|-----|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|-----|
| BAL | | | | | | | | | | | | | |
| BLU | 0,000801 | | | | | | | | | | | | |
| CUT | 0,201441 | 0,088417 | | | | | | | | | | | |
| KAM | 0,164496 | 0,156476 | 0,171574 | | | | | | | | | | |
| LAN | 0,107052 | 0,164271 | 0,060774 | 0,154542 | | | | | | | | | |
| LUN | 0,079577 | 0,076468 | 0,067809 | 0,097922 | 0,117718 | | | | | | | | |
| MAN | 0,12655 | 0,055412 | 0,162027 | 0,07891 | 0,073448 | 0,106756 | | | | | | | |
| MUL | 0,13998 | 0,153908 | 0,103981 | 0,131695 | 0,071607 | 0,172205 | 0,107984 | | | | | | |
| OBR | 0,112019 | 0,084681 | 0,098626 | 0,106568 | 0,216642 | 0,159714 | 0,045017 | 0,065042 | | | | | |
| PLA | 0,139618 | 0,11118 | 0,082283 | 0,097426 | 0,179182 | 0,155014 | 0,077423 | 0,050775 | 0,072719 | | | | |
| SIL | 0,016175 | 0,191983 | 0,156895 | 0,08078 | 0,047379 | 0,085416 | 0,139032 | 0,110211 | 0,108893 | 0,147869 | | | |
| SMO | 0,215048 | 0,089054 | 0,066703 | 0,090073 | 0,15975 | 0,093084 | 0,089633 | 0,130803 | 0,1268 | 0,11043 | 0,068103 | | |
| VRU | 0,193178 | 0,082158 | 0,093824 | 0,12952 | 0,168743 | 0,06866 | 0,052086 | 0,116784 | 0,042168 | 0,083203 | 0,082285 | 0,073752 | |

Ln (distance):

| | BAL | BLU | CUT | KAM | LAN | LUN | MAN | MUL | OBR | PLA | SIL | SMO | VRU |
|-----|---------|---------|---------|---------|---------|---------|---------|---------|-----|-----|-----|-----|-----|
| BAL | | | | | | | | | | | | | |
| BLU | 2,30259 | | | | | | | | | | | | |
| CUT | 11,71 | 9,86246 | | | | | | | | | | | |
| KAM | 11,7099 | 12,2818 | 12,0995 | | | | | | | | | | |
| LAN | 11,6082 | 12,2861 | 10,7962 | 11,6397 | | | | | | | | | |
| LUN | 11,6081 | 6,98193 | 10,8162 | 8,65434 | 11,1055 | | | | | | | | |
| MAN | 12,3314 | 6,97261 | 12,0303 | 11,1184 | 8,45468 | 10,4038 | | | | | | | |
| MUL | 11,6159 | 11,7196 | 11,0354 | 11,1196 | 11,1861 | 11,9189 | 12,1014 | | | | | | |
| OBR | 11,6158 | 11,5997 | 8,44484 | 11,1195 | 11,9087 | 12,2062 | 10,7657 | 11,1476 | | | | | |

| | | | | | | | | | | | | | |
|-----|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|--|
| PLA | 12,3348 | 11,6074 | 8,44269 | 12,1122 | 11,9086 | 11,8765 | 10,7869 | 7,99294 | 10,4834 | | | | |
| SIL | 7,13728 | 11,6897 | 11,672 | 10,7038 | 10,4529 | 12,2309 | 12,0367 | 12,2213 | 12,3594 | 11,8178 | | | |
| SMO | 11,6797 | 11,0467 | 11,6495 | 10,7265 | 12,4249 | 11,0776 | 11,0633 | 11,8101 | 11,806 | 12,1283 | 12,1532 | | |
| VRU | 11,6796 | 11,0465 | 11,6569 | 12,0564 | 12,4237 | 11,0775 | 7,5689 | 11,81 | 9,70021 | 11,7755 | 10,2103 | 12,1378 | |

distances (m):

| | BAL | BLU | CUT | KAM | LAN | LUN | MAN | MUL | OBR | PLA | SIL | SMO | VRU |
|-----|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|-----|
| BAL | | | | | | | | | | | | | |
| BLU | 10 | | | | | | | | | | | | |
| CUT | 121782 | 121772 | | | | | | | | | | | |
| KAM | 109997 | 109987 | 226704 | | | | | | | | | | |
| LAN | 110844 | 110834 | 227482 | 1258 | | | | | | | | | |
| LUN | 118147 | 118137 | 19196 | 215740 | 216656 | | | | | | | | |
| MAN | 1077 | 1067 | 122959 | 109067 | 109906 | 119340 | | | | | | | |
| MUL | 62736 | 62726 | 179786 | 48834 | 49820 | 167756 | 62031 | | | | | | |
| OBR | 4651 | 4641 | 117241 | 114635 | 115480 | 113519 | 5735 | 67401 | | | | | |
| PLA | 67483 | 67473 | 182088 | 44526 | 45547 | 172200 | 66535 | 4697 | 72118 | | | | |
| SIL | 148548 | 148538 | 34644 | 248916 | 248617 | 32985 | 150072 | 200025 | 143846 | 205023 | | | |
| SMO | 64706 | 64696 | 180123 | 47369 | 48381 | 168832 | 63789 | 1937 | 69400 | 2960 | 203059 | | |
| VRU | 134601 | 134591 | 35717 | 233149 | 134053 | 16321 | 135651 | 185035 | 130021 | 189705 | 27182 | 186803 | |

Mantel test, 10000 permutations

(statistic: Spearman Rank correlation coefficient)

Test of isolation by distance (One tailed Pvalue):


Pr (correlation > observed correlation) =0.0002 under null hypothesis

Other one tailed Pvalue:

Pr (correlation < observed correlation) =0.9998 under null hypothesis

Appendix 5

a Adriatic Sea coastline and topography, included from Artegiani *et al.* (1996)



b Seasonal maps of surface mean flow in the Adriatic basin, included from Poulain (2001)

